

## Acute Hyponatremia Induced by Bowel Preparations for Colonoscopy: Identification of Patients at Risk

Eli D Ehrenpreis<sup>1\*</sup>  
Nyi Nyi Tun<sup>2</sup>  
Dino Melitas<sup>3</sup>

<sup>1</sup>Adjunct Professor of Pediatric Gastroenterology, University of Miami Miller School of Medicine, Professor of Medicine, Rosalind Franklin University Medical School, and Associate Director for Research, Advocate Lutheran General Hospital, Park Ridge, IL

<sup>2</sup>Department of Medicine, Miami Valley Hospital, Dayton OH

<sup>3</sup>Gastroenterology Division, Providence-Providence Park Hospital, Southfield, MI

## Abstract

**Introduction:** Iatrogenic acute hyponatremia has been observed following bowel preparation for colonoscopy and is believed to be associated with hypo-osmotic preparations.

**Aim:** To review cases of hyponatremia attributed to bowel preparations for colonoscopy reported to the Food and Drug Administration (FDA) Adverse Event Reporting System (FAERS).

**Methods:** All adverse event (ADR) reports to the FAERS reported between March 2008 and May 2014 were reviewed. The data was filtered to cases using the term "colonoscopy" with primary ADRs of "hyponatremia" and/or seizure". Bowel preparations reviewed (including alternative spellings of brand names) included: *OsmoPrep™*, *MoviPrep™*, *SuPrep™*, *Visicol™*, *Magnesium citrate*, *Golytely™*, *MiraLAX™*, *Trilyte™*, *Gavilyte™*, and *Colyte™*. Complete reports of all cases were obtained from the FDA using the Freedom of Information Act (FOIA).

**Results:** We reviewed 40 cases of colonoscopy preparations with primary diagnosis of hyponatremia; 15 hyponatremia cases also reported seizures. We also reviewed 17 cases with primary ADRs of seizures without hyponatremia. Serum sodium levels were available in 27 of 40 hyponatremia cases with a mean value of 118 mEq/L. The average serum sodium level in hyponatremic patients developing a seizure was 115 mEq/L. *OsmoPrep™*, *MoviPrep™* and *SuPrep™* accounted for the majority of hyponatremia cases (35%, 32.5% and 10% respectively). Two patients with hyponatremia died from seizures followed by cerebral edema. Predisposing factors for hyponatremia induced by colonoscopy preparations include female gender (72.5%); excess fluid ingestion (17.5%), hypothyroidism (15%), thiazide diuretics use (10%), SSRI use (7.5%) and combination of more than one preparation (5%).

**Conclusions:** Acute hyponatremia is an uncommon but serious complication of a variety of colonoscopy bowel preparations. Identification of patients at risk, avoiding combinations of preparations and limiting excess fluid intake are strategies to prevent iatrogenic hyponatremia from colonoscopy preparation.

## Introduction

Colonoscopy is a common procedure utilized for both diagnostic and therapeutic purposes in daily clinical settings. According to the data from the Centers for Disease Control and Prevention (CDC), more than 10 million colonoscopies are performed annually the U.S. alone [1]. The overall risk of serious complications following colonoscopy is low. In a review of 12 studies with 57,742 screening colonoscopies, serious harm occurred in 2.8 per 1,000 examinations and over 85 % of the complications occurred in the setting of polypectomy [2].

An excellent bowel preparation is essential for diagnostic accuracy and safety of the colonoscopy procedure [3,4]. Bowel preparations should empty the entire colon of all fecal material without any damage of the colonic mucosa. The ideal bowel preparation should be effective, tolerable, inexpensive and convenient to use [3-6]. Bowel preparations should also not cause discomfort or fluid and electrolytes imbalance. The preparation agents may be isosmotic (e.g. Polyethylene glycol (PEG) preparation), hyposmotic (e.g. a combination of PEG-3350 and sports drink), hyperosmotic (e.g. sodium sulfate, magnesium citrate, sodium phosphate) or a combination of iso- and hyper-osmotic (e.g. combination of sodium picosulfate and magnesium citrate). Our group reported fluid and electrolytes shifts occurring after hyperosmotic colonoscopy preparations shortly after their acceptance on the market [7]. Bowel regimens are also variable in regard to ingested fluid volume and include high volume (4 liters/128 oz) or low volume (2 liters/64 oz) preparations. Not all of the regimens are approved for use by the Food and Drug Administration (FDA) [6]. Over-the-counter preparation regimens involving mixing with sports drinks, *MiraLAX™* +

## Article Information

**DOI:** 10.31021/dddj.20181104  
**Article Type:** Research Article  
**Journal Type:** Open Access  
**Volume:** 1 **Issue:** 1  
**Manuscript ID:** DDDDJ-1-104  
**Publisher:** Boffin Access Limited  
**Received Date:** 15 April 2018  
**Accepted Date:** 10 May 2018  
**Published Date:** 22 May 2018

### \*Corresponding author:

#### Eli D Ehrenpreis

Adjunct Professor of Pediatric Gastroenterology  
University of Miami Miller School of Medicine  
Professor of Medicine  
Rosalind Franklin University Medical School and  
Associate Director for Research  
Advocate Lutheran General Hospital  
Park Ridge, IL.  
USA  
E-mail: e2bioconsults@gmail.com

**Citation:** Ehrenpreis ED, Tun NN, Melitas D. Acute Hyponatremia Induced by Bowel Preparations for Colonoscopy: Identification of Patients at Risk. Drug Des Dev Deliv J. 2018 May;1(1):104

**Copyright:** © 2018 Ehrenpreis ED, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 international License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. Vium mor li, poracidisus es condam

*Gatorade* combination, are hyposmolar regimens not approved by the FDA [8]. Concerns regarding with the risk of hyponatremia from these preparations have been voiced in the medical literature [9-11]. A 2001 study published in *The Lancet* reported an incidence (7.5%) of hyponatremia after colonoscopy [12]. The administration of bowel regimens can predispose to the development of hyponatremia by a number of mechanisms which result in relative increase in the ratio of body water to sodium. Acute hyponatremia can produce detrimental neurological complications culminating in seizures, coma and death [13].

We reviewed cases of bowel preparations with the occurrence of hyponatremia and/or seizures reported to the Food and Drug Adverse Event Reporting System (FAERS). We focused on the characteristics that may predispose to the development of hyponatremia.

## Materials and Methods

The FAERS is a publicly accessible database, used for post-marketing safety surveillance for all FDA-approved drugs and therapeutic agents. The adverse events are voluntarily reported by physicians, pharmacists and other healthcare professionals as well as drug manufacturers, patients, and attorneys. The database was searched for cases of bowel preparation associated with the term "colonoscopy" with reports of a primary ADR diagnosis of "hyponatremia" or a primary diagnosis of "seizure". The individual files were downloaded for the period between March 2008 and May 2014. The types of bowel preparations utilized in the reported cases included *OsmoPrep™*, *MoviPrep™*, *SuPrep™*, *Visicol™*, *Magnesium citrate*, *Golytely™*, *MiraLAX™*, *Trilyte™*, *Gavilyte™*, and *Colyte™*. Complete reports of all cases from the FAERS database were obtained from the FDA using the Freedom of Information Act (FOIA). The cases were reviewed individually and analyzed to establish authenticity and to remove duplicated cases. All the reports underwent careful examination to confirm the occurrence of hyponatremia and/or seizures after colonoscopy bowel preparation. The reports were further evaluated and the information was gathered on age, gender, type of bowel preparation implicated, hyponatremia, other reported electrolytes abnormalities, potential contributing factors for hyponatremia and underlying disease states. Furthermore, serum sodium levels were collected if reported and the neurologic complications such as seizures were recorded. Concurrent medications including Selective Serotonin Inhibitors (SSRIs) and thiazides diuretics were also recorded.

## Results

We reviewed 40 cases that were reported to have a primary diagnosis of the ADR of hyponatremia and 17 cases reported to develop seizures without hyponatremia. Fifteen (37.5%) of hyponatremia cases also developed seizures. None of these patients were reported to have hyponatremia prior to the use of the bowel preparation. Serum sodium levels were available in 27 of 40 hyponatremia cases and ranged from 103 to 128 mEq/L (normal serum sodium = 135-145 mEq/L) with a mean value of 118 mEq/L. The average serum sodium level in hyponatremic patients developing a seizure was 115 mEq/L (Figure 1). The most common reported preparations associated with hyponatremia were *OsmoPrep™* (14 cases, 35%), *MoviPrep™* (13 cases, 32.5%) and *SuPrep™* (4 cases, 10%), (Table 1).

Two patients with hyponatremia following colonoscopy preparation died from cerebral edema and seizures. Both of these patients were found to have ingested *MoviPrep™*.

The patient characteristics of the hyponatremia cases are shown in Table 2. The factors that were associated with the development of hyponatremia included female gender (72.5%), fluid ingestion in excess of recommended amounts (17.5%), hypothyroidism (15%), thiazide diuretics use (10%), use of selective serotonin reuptake Inhibitors (SSRI) (7.5%), cirrhosis (5%) and the combined use of more than one bowel preparation (5%). Only 2 patients with primary reports of seizure without hyponatremia following colonoscopy preparation had prior history of seizure disorder.

## Discussion

Bowel preparation agents using a variety of regimens are generally regarded as safe and well tolerated if the agents are appropriately prescribed and the preparation instructions are correctly followed [14-17]. Despite their overall safety profile, there are known gastrointestinal (GI) and non-GI related adverse events associated with their use. GI complications include nausea, vomiting, abdominal distension, and ileus. Other known complications include electrolyte abnormalities including hyponatremia, seizures, encephalopathy, exacerbation of congestive heart failure and pancreatitis. Phosphate nephropathy followed by chronic renal failure is uniquely associated with the use of sodium phosphate containing colonoscopy preparations and has limited their use [18-26].

Our group has determined that renal adverse drug reactions (phosphate nephropathy, renal failure) from sodium phosphate-based colonoscopy preparation are more common in women. Lower body weight is another risk factor for kidney injury from sodium phosphate-based colonoscopy preparations [27].

The current study of ADR reports to the FAERS describes the occurrence of hyponatremia secondary to a variety of bowel

Bowel preparation	Primary Diagnosis	
	Hyponatremia (n=40)	Seizures without Hyponatremia (n=17)
<i>OsmoPrep™</i>	14 (35%)	3 (17.6%)
<i>MoviPrep™</i>	13 (32.5%)	5 (29.4%)
<i>SuPrep™</i>	4 (10%)	1 (5.9%)
<i>Visicol™</i>	1 (2.5%)	3 (17.6%)
Magnesium citrate	1 (2.5%)	0
<i>Golytely™</i>	1 (2.5%)	2 (11.8%)
<i>MiraLAX™</i>	4 (10%)	2 (11.8%)
<i>Trilyte™</i>	1 (2.5%)	0
<i>Gavilyte™</i>	0	1 (5.9%)
<i>Colyte™</i>	1 (2.5%)	0

**Table 1:** Cases of Primary Diagnosis of Hyponatremia and Cases of Primary Diagnosis of Seizure without Hyponatremia According to Bowel Preparation Regimens

Gender	Female – 29 (72.5%), Male – 10 (25%), Unreported – 1 (2.5%)
Age group	30-39 Yr – 3 (7.5%), 40-49 Yr – 2 (5%), 50-59 Yr – 14 (35%), 60-69 Yr – 12 (30%), 70-79 Yr – 2 (5%), 80 Yr and above – 1 (2.5%), Unreported age – 6 (15%)
Other electrolytes abnormalities	Hypokalemia – 20 (50%), Hypocalcemia – 6 (15%)
Thiazide diuretics	4 (10%)
SSRI	3 (7.5%)
Excess fluid ingestion*	7 (17.5%)
Hypothyroidism	6 (15%)
Mixed use of more than one bowel preparation*	2 (5%)
Cirrhosis	2 (5%)

**Table 2:** Characteristics of patients who developed hyponatremia secondary to bowel preparation (n=40)

\*3 cases of *MoviPrep™*, 2 cases of *OsmoPrep™*, 1 case of *Visicol™*, 1 case of *SuPrep™*

\*One reported case used a combination of *MiraLAX™* and bisacodyl and the other used both *MoviPrep™* and magnesium citrate.




Figure 1: Hyponatremic patients with reported serum sodium level

regimens. In our study, *OsmoPrep™* and *MoviPrep™* accounted for nearly 70% of hyponatremia cases. *OsmoPrep™* is a low volume hyperosmotic regimen containing monobasic and dibasic sodium phosphate compounds whereas *MoviPrep™* is a low volume isosmotic regimen composed of Polyethylene Glycol (PEG) and ascorbic acid. The latter component of *MoviPrep™* also contributes to the cathartic effect of the preparation [17].

As anticipated, 10% of cases were described with the use of *SuPrep™*, a hypo-osmolar preparation. A publication issued by UK National Patient Safety Agency (NPSA) reported 218 patient safety incidents associated with the use of bowel cleansing preparations over a 5-year period. 6% of the 218 incidents caused moderate patient harm and one patient death was reported. These data are derived from a voluntary reporting system, similar to FAERS in the U.S. Hyponatremia was a significant adverse effect highlighted in the NPSA report [17]. An observational study was undertaken to assess the incidence of hyponatraemia in colonoscopy patients then published in 2001 by The Lancet. In the study, 40 patients (14 women, 26 men, mean age 57.4 years) were assessed before and after colonoscopy with 20 gastroscopy patients (10 women, 10 men, mean age 55.8 years) serving as controls. On the morning of procedure, the colonoscopy patients drank 2-3 liters of bowel preparation solution containing PEG and balanced electrolytes. Serum levels of sodium and arginine vasopressin (ADH) were collected in all patients before and after the procedure. The study found 3 (7.5%) of 40 colonoscopy patients had a serum sodium concentration of 130 mEq/L or lower in association with elevated ADH levels. By contrast, gastroscopy patients in the control group developed neither hyponatremia nor elevated ADH levels [12]. Except for this small study in The Lancet, the incidence of hyponatremia in after bowel preparation for colonoscopy has not been reported, as most of the literature is based on individual case reports. The methodology employed in data collection using the FAERS also does not allow for determination of the incidence of this condition, due to the voluntary nature of ADR reports to the FDA [27].

The association of hyponatremia with individual colonoscopy preparations can also not be determined using this method. Nonetheless, our study emphasizes that hyponatremia may occur following a variety of colonoscopy preparations and is not limited to hypo-osmolar preparations.

PEG is a non-absorbable agent that passes through the bowel without significant net absorption or secretion and therefore typically does not result in fluid and electrolyte shifts [6]. The total net gain or loss of water and sodium from bowel clearance with PEG is unlikely to exceed 700 ml or 30 mmol respectively and therefore the clinical impact of those changes are minimal [4]. In

contrast, sodium phosphate regimens are hyperosmotic promoting pronounced fluid and electrolytes shifts and colonic evacuation by drawing large amount of water (1-1.8 liters of water per 45 ml of preparation) [28]. A pharmacokinetic study of sodium phosphate based colon cleansing solution found elevated serum phosphate levels (mean serum phosphate at 120 min was  $7.8 \pm 0.5$  mg/dL) in smaller individuals (less than 55 kg) after ingestion of the solution [29]. Hyperosmotic colonoscopy preparations, especially those containing sodium phosphate should generally be avoided and are contraindicated in patients with chronic kidney disease, congestive cardiac failure, cirrhosis [28].

PEG-based isosmotic bowel regimens can be either high volume (4L) preparations such as PEG mixed with balanced electrolyte solution (PEG-ELS) or newer low volume (2L) preparations such as *MoviPrep™*, *Halflytely™* or *MiraLAX™*. The latter two are usually co-administered with bisacodyl. Bisacodyl used as an adjunct to PEG-based preparations, has been reported to be associated with abdominal cramping and ischemic colitis [30-31].

Although PEG-ELS are generally well tolerated, 5% to 15% of patients do not complete the preparation because of poor palatability and/or large volume [6-14]. Accordingly, low volume PEG based regimens are more desirable for patients who cannot tolerate high volume preparations.

However, patients who opt for low volume preparations tend to drink more water or clear liquids than instructed. According to one European study, it is found that 82.1% of patients who took PEG + ascorbic acid regimen drank at least 1 L or more additional water or clear liquids, while 66.1% of patients who took sodium phosphate regimen drank 2 L additional water or clear liquids [32]. Not all of the ingested water stays within the bowel lumen and part of it will be absorbed and therefore overzealous drinking of water can lead to water intoxication and hyponatremia [17]. This finding is confirmed in our study. We demonstrated that 7 of 40 hyponatremia cases, (17.5%), drank additional water or hypotonic fluids beyond the recommended amounts. A new finding in our study is the risk of hyponatremia is increased when low volume preparations are combined with other preparations.

Concerns about iatrogenic hyponatremia from the use of over-the-counter PEG-based hyposmolar regimens have been cited in the medical literature. A common form of these regimens is *MiraLAX™* + Gatorade combination (238 grams of PEG-3350 without electrolytes mixed in 64 ounces of Gatorade) which is not FDA approved. This preparation has achieved popularity because of lower volume and improved palatability compared to standard high volume PEG and electrolyte solution regimen as well as cost saving [8]. However, *MiraLAX™* + *Gatorade* regimen is not an osmotically balanced

solution and its safety has not been evaluated in sufficiently large studies [9]. A recent case series identified severe hyponatremia and serious adverse events, including cardiac arrhythmias and ICU admissions in 14 low-risk patients who used *MiraLAX™ + Gatorade* bowel preparations [10]. In addition, a study based on an endoscopic database of over 8,400 patients found a higher incidence of severe hyponatremia with *MiraLAX™ + Gatorade* versus osmotically balanced PEG bowel preparations [11].

Hyponatremia is commonly defined as a serum sodium concentration below 135 mEq/L [33-34]. Hyponatremic state with serum sodium level above 130 mEq/L usually does not cause any clinical symptoms. The symptoms of hyponatremia typically manifest when there is acute and marked reduction in the serum sodium concentration. Nausea and malaise, which are the earliest manifestations, may be seen when the serum sodium concentration falls below 125 to 130 mEq/L. If the serum sodium concentration falls below 115 to 120 mEq/L and further, headache, lethargy, obtundation, seizures, encephalopathy, cerebral edema, stupor, coma and respiratory arrest will ensue [13,35,36]. Bowel regimens may cause hyponatremia via several overlapping mechanisms. One of these mechanisms is the production of osmotic diarrhea. This causes volume depletion and high and low pressure baroreceptor-mediated nonosmotic stimulation of ADH secretion. The end result of high ADH state is free water retention resulting in hyponatremia [37]. Other mechanisms of bowel cleansing agents-induced hyponatremia include overenthusiastic consumption of large amount of water, concomitant parenteral hypotonic fluid administration, renal failure and increased ADH-release states induced by several non-osmotic stimuli including pain, nausea, and abdominal manipulation related to the colonoscopy procedure [12,17,38]. Our study raises the possibility that concomitant medications such as thiazide diuretics and SSRIs may also contribute to the development of hyponatremia. Thiazide-induced hyponatremia has an underlying complex mechanism and is thought to result from the combination of sodium loss induced by the diuretic and enhanced water re-absorption due to both ADH-dependent and ADH-independent pathways [39,40]. Various SSRIs can predispose patients to develop hyponatremia by causing SIADH [41]. An additional new finding in our study is that hyponatremia from colonoscopy preparations is a more common occurrence in female patients, as 72.5% of cases reported to the FAERS occurred in females. Hyperphosphatemia and calcium-phosphate nephropathy from sodium phosphate containing colonoscopy preparations are also more common in females. This has been attributed to lower overall body weight and smaller amount of total body water and vascular volume in females [27,29]. Pre-existing co-morbid states such as hypothyroidism and cirrhosis of the liver appeared as contributing factors to the occurrence of hyponatremia in our study group. Although incompletely understood, hypothyroidism-induced hyponatremia is thought to be related to decreased ability to excrete free water [42,43]. A variety of factors are implicated in the predisposition of cirrhotic patients to develop hyponatremia and the most important factor is diminished effective circulatory volume and arterial vasodilatation [44,45].

The occurrence of seizures occurring in patients undergoing colonoscopy and attributed to colonoscopy preparations has not received attention in the medical literature. Reports to the FAERS appear to implicate a variety of colonoscopy preparations in this form of ADR (Table 1). The mechanism of seizure development in colonoscopy patients without hyponatremia is unclear and possibly multifactorial. Colonoscopy is a procedure involving the use of pharyngeal anesthesia and sedation [46]. Lidocaine and fentanyl are frequently used agents for colonoscopy and they can provoke seizures. Patients may be on concomitant medications such as antibiotics, antidepressants and antipsychotics which can also provoke seizures [47]. Concurrent acute medical illnesses and metabolic disturbances such as hypoglycemia, hyperglycemia, cerebral vascular accident, cerebral anoxia, hypocalcemia, hypomagnesemia may also provoke seizures in colonoscopy patients without hyponatremia [48,49]. Further elucidation and research is needed to better understand the pathogenesis of seizures in that subset of patient population.

In summary, the exact incidence of iatrogenic hyponatremia induced by bowel preparation is unknown, although the condition appears to be rare based on the small number of reports to FAERS. Physicians should be aware of its occurrence and complications. Acute hyponatremia can result in neurologic damage and even death. Female patients appear to be at higher risk. Written instructions should include avoidance of excess free water intake. Combination of more than one colonoscopy preparations may also represent a risk for the development of hyponatremia and should be avoided.

## References

1. Seeff LC, Richards TB, Shapiro JA, Nadel MR, Manninen DL, et al. How many endoscopies are performed for colorectal cancer screening? Results for the CDC's survey of endoscopic capacity. *Gastroenterology*. 2004 Dec;127(6):1670-1677.
2. Whitlock EP, Lin JS, Liles E, Beil TL, Fu R. Screening for colorectal cancer: a targeted, updated systematic review for the U.S. Preventive Services Task Force. *Ann Intern Med*. 2008 Nov; 149(9):638.
3. Froehlich F, Wietlisbach V, Gonvers JJ, Burnand B, Vader JP. Impact of colonic cleansing on quality and diagnostic yield of colonoscopy: the European Panel of Appropriateness of Gastrointestinal Endoscopy European multicenter study. *Gastrointest Endosc*. 2005 Mar;61(3):378-384.
4. Belsey J, Epstein O, Heresbach D. Systematic review: adverse event reports for oral sodium phosphate and polyethylene glycol. *Aliment Pharmacol Ther*. 2009 Jan;29(1):15-28.
5. DiPalma JA, Brady CE. Colon cleansing for diagnostic and surgical procedures: polyethylene glycol-electrolyte lavage solution. *Am J Gastroenterol* 1989 Sep;84(9):1008-1016.
6. ASGE Standards of Practice Committee, Saltzman JR, Cash BD, Pasha SF, Early DS, Muthusamy VR et al. Bowel preparation before colonoscopy. *Gastrointestinal Endoscopy*. 2015 Apr;81(4):781-794.
7. Ehrenpreis ED, Noguerras JJ, Botoman VA, Bonner GF, Zaitman D, et al. Serum electrolyte abnormalities secondary to Fleet's Phospho-Soda colonoscopy prep: A review of three cases. *Surg Endosc*. 1996 Oct;10(10):1022-1024.
8. Frederick K. Shieh, Naresh Gunaratnam, Sagal O. Mohamud, Schoenfeld P. *MiraLAX-Gatorade Bowel Prep vs. GoLytey Prior to Screening Colonoscopy: An Endoscopic Database Study in a Community Hospital*. *J Clin Gastroenterol*. 2012 Nov-Dec;46(10):e96-e100.
9. Schoenfeld P. Safety of MiraLAX/Gatorade bowel preparation has not been established in appropriately designed studies. *Clin Gastroenterol Hepatol*. 2013 May;11(5):582.
10. Lewis J, Schoenfeld P. Severe Hyponatremia and Miralax-Gatorade Bowel Preparation. *Am J of Gastroenterol*. 2011 Jan;106(Suppl 2):S582.
11. Schoenfeld P, Elliott E. Increased Risk of Severe Hyponatremia with Miralax-Gatorade vs Standard Bowel Preparation. *Am J of Gastroenterology*. 2011;106(Suppl 2):S583.
12. Cohen CD, Keunike C, Schiemann U, Schroppel B, Siegert S, et al. Hyponatremia as a complication of colonoscopy. *Lancet*. 2001 Jan;357(9252):282-283.
13. Sterns RH. Disorders of plasma sodium--causes, consequences, and correction. *N Engl J Med*. 2015 Jan;372(1):55-65.
14. Wexner SD, Beck DE, Baron TH, Fanelli RD, Hyman N, et al. A consensus document on bowel preparation before colonoscopy: prepared by a task force from the American Society of Colon and Rectal Surgeons (ASCRS), the American Society for Gastrointestinal Endoscopy (ASGE), and the Society of American Gastrointestinal and Endoscopic Surgeons (SAGES). *Gastrointest Endosc*. 2006 Jun;63(7):894-909.
15. Vanner SJ, MacDonald PH, Paterson WG, et al. A randomized prospective trial comparing oral sodium phosphate with standard polyethylene glycol-based lavage solution (Golytey) in

- the preparation of patients for colonoscopy. *Am J Gastroenterol*. 1990Apr;85(4):422-447.
16. Reddy DN, Rao GV, Sriram PV. Efficacy and safety of oral sodium phosphate versus polyethylene glycol solution for bowel preparation for colonoscopy. *Indian J Gastroenterol*. 2002Nov-Dec;21(6):219-221.
  17. Connor A, Tolan D, Hughes S, Carr N, Tomson C. Consensus guidelines for the safe prescription and administration of oral bowel-cleansing agents. *Gut*. 2012 Nov;61(11):1525-1532.
  18. Granberry MC, White LM, Gardner SF. Exacerbation of congestive heart failure after administration of polyethylene glycol-electrolyte lavage solution. *Ann Pharmacother*. 1995 Dec;29(12):1232-1235.
  19. Nagler J, Poppers D, Turetz M. Severe hyponatremia and seizure following a polyethylene glycol-based bowel preparation for colonoscopy. *J Clin Gastroenterol*. 2006 Jul;40(6):558-559.
  20. Frizelle FA, Colls BM. Hyponatremia and seizures after bowel preparation: report of three cases. *Dis Colon Rectum*. 2005 Feb;48(2):393-396.
  21. Franga DL, Harris JA. Polyethylene glycol-induced pancreatitis. *Gastrointest Endosc*. 2000 Dec;52(6):789-791.
  22. Mackey AC, Schaffer D, Prizant R. Seizure associated with the use of visicol for colonoscopy. *N Engl J Med*. 2002 Jun;346(26):2095.
  23. Rose M, Jacob LS. Seizure associated with the use of visicol for colonoscopy. *N Engl J Med*. 2002 Jul;347(4):295-296.
  24. Schroppel B, Segerer S, Keuneke C, Cohen CD, Schlöndorff D, et al. Hyponatremic encephalopathy after preparation for colonoscopy. *Gastrointest Endosc*. 2001Apr;53(4):527-529.
  25. Markowitz GS, Stokes MB, Radhakrishnan J, D'Agati VD. Acute phosphate nephropathy following oral sodium phosphate bowel purgative: an underrecognized cause of chronic renal failure. *Am Soc Nephrol*. 2005 Sep;16(11):3389-3396.
  26. Heher EC, Thier SO, Rennke H, Humphreys BD. Adverse renal and metabolic effects associated with oral sodium phosphate bowel preparation. *Clin J Am Soc Nephrol*. 2008 Sep;3(5):1494-1503.
  27. Ehrenpreis ED, Parakkal D, Semer R, Du H. Renal risks of sodium phosphate tablets for colonoscopy preparation: a review of adverse drug reactions reported to the US Food and Drug Administration. *Colorectal Dis*. 2011 Sep;13(9):e270-275.
  28. Schiller LR. Clinical pharmacology and use of laxatives and lavage solutions. *J Clin Gastroenterol*. 1999 Jan;28(1):11-18.
  29. Ehrenpreis ED. Increased serum phosphate levels and calcium fluxes are seen in smaller individuals after a single dose of sodium phosphate colon cleansing solution: a pharmacokinetic analysis. *Aliment Pharmacol Ther*. 2009 Jun;29(11):1202-1211.
  30. Baudet JS, Castro V, Redondo I. Recurrent ischemic colitis induced by colonoscopy bowel lavage. *Am J Gastroenterol*. 2010 Mar;105(3):700-701.
  31. Lopez Morra HA, Fine SN, Dickstein G. Colonic ischemia with laxative use in young adults. *Am J Gastroenterol*. 2005 Sep;100(9):2134-2136.
  32. Bitoun A, Ponchon T, Barthet M, Coffin B, Dugué C. Results of a prospective randomised multicentre controlled trial comparing a new 2-L ascorbic acid plus polyethylene glycol and electrolyte solution vs. sodium phosphate solution in patients undergoing elective colonoscopy. *Aliment Pharmacol Ther*. 2006 Dec;24(11-12):1631-1642.
  33. Adrogué HJ, Madias NE. Hyponatremia. *N Engl J Med*. 2000 May;342(21):1581-1589.
  34. Spasovski G, Vanholder R, Alolio B, Annane D, Ball S, et al. Clinical practice guideline on diagnosis and treatment of hyponatraemia. *Nephrol Dial Transplant*. 2014 Apr;29.
  35. Ellis SJ. Severe hyponatraemia: complications and treatment. *QJM*. 1995;88(12):905.
  36. Moritz ML, Ayus JC. The pathophysiology and treatment of hyponatraemic encephalopathy: an update. *Nephrol Dial Transplant*. 2003 Dec;18(12):2486-2491.
  37. Anderson RJ, Chung HM, Kluge R, Schrier RW. Hyponatremia: a prospective analysis of its epidemiology and the pathogenetic role of vasopressin. *Ann Intern Med*. 1985 Feb;102(2):164-168.
  38. Cohen LB, Kastenber DM, Mount DB, Safdi AV. Current Issues in Optimal Bowel Preparation: Excerpts From a Roundtable Discussion Among Colon-Cleansing Experts. *Gastroenterol Hepatol (N Y)*. 2009 Nov;5(11 Suppl 19): 3-11.
  39. Ashraf N, Locksley R, Arief AI. Thiazide-induced hyponatremia associated with death or neurologic damage in outpatients. *Am J Med*. 1981;70(6):1163.
  40. Cesar KR, Magaldi AJ. Thiazide induces water absorption in the inner medullary collecting duct of normal and Brattleboro rats. *Am J Physiol*. 1999 Nov;277(5 Pt 2):F756.
  41. Fabian TJ, Amico JA, Kroboth PD, Mulsant BH, Corey SE, et al. Paroxetine-induced hyponatremia in older adults: a 12-week prospective study. *Arch Intern Med*. 2004;164(3):327-332.
  42. Schrier RW. Body water homeostasis: clinical disorders of urinary dilution and concentration. *J Am Soc Nephrol*. 2006;17(7):1820.
  43. Derubertis FR Jr, Michelis MF, Bloom ME, Mintz DH, Field JB, Davis BB. Impaired water excretion in myxedema. *Am J Med*. 1971;51(1):41.
  44. Abelmann WH. Hyperdynamic circulation in cirrhosis: a historical perspective. *Hepatology*. 1994;20(5):1356.
  45. Groszmann RJ. Hyperdynamic circulation of liver disease 40 years later: pathophysiology and clinical consequences. *Hepatology*. 1994;20(5):1359.
  46. Cohen LB, DeLegge MH, Aisenberg J, et al. AGA institute review of endoscopic sedation. *Gastroenterology*. 2007;133:675.
  47. Eisenschenk S, Gilmore RL. Seizures associated with nonneurologic medical conditions. In: *The Treatment of Epilepsy: Principles and Practice*, 4th ed, Gupta A, Lachhwani DK (Eds), Lippincott Williams & Wilkins, Philadelphia 2006.
  48. Beghi E, Carpio A, Forsgren L, Hesdorffer DC, Malmgren K, et al. Recommendation for a definition of acute symptomatic seizure. *Epilepsia*. 2010;51(4):671.
  49. Riggs JE. Neurologic manifestations of electrolyte disturbances. *Neurol Clin*. 2002;20(1):227.