

Mycoflora of Ground Water Associated with Al-Teeb well in Iraq

Hussein Al-Nasrawi

Assistant professor, Southern Technical University, Iraq

Abstract

Al-Teeb well is an important groundwater source for drinking water used by marsh dwellers in southern area of Iraq. The mentioned water facility supply more than 250,000 marsh dwellers of Misan province in Iraq with drinking water by tanker trucks. 43 fungi were isolated from sediments of Al-Teeb well, the isolated fungi included one new first record species *Allysedium sp.*, added to Iraqi fungi, mentioned new record fungus was described and illustrated, has been kept in Basrah herbarium under no. 20101.

Keywords

Groundwater; Fungi; Sediments; Iraq

Introduction

Bacteria and fungi are the most common microorganisms of soil; however molds outnumber bacteria, under conditions of poor aeration, low temperature, and acidity, because they tolerate these conditions more easily. Molds are types of fungi distinguish by their mycelia; threads grow in different environments specially soil and aquatic ecosystems [1].

In aquatic ecosystem fungi consider one of microorganisms that play an important role in biodegradation process of cellulose containing materials and decompose undesired organic materials submerged in water and contribute in food chain. The potential of using white-rot fungi for the degradation of pollutants has been extensively researched in-vitro and has mainly been attributed to their ability to degrade lignin [2].

Fungi in water distribution systems are well known and have gained importance; it is unsurprising that fungi are isolated from the surface or underground raw water in reservoirs and distribution systems, since they are found in almost every environmental niche [3].

Most studies involved in evaluating the potential of white-rot fungi in bio-remediation have been done under lab conditions. However, non-white rot fungi will not fully mineralize of PAH, whereas white rot fungi are capable of doing so by introducing oxygen into the rings, rendering the molecules more soluble and chemically reactive. This reaction is produced via extracellular enzymes produced by white-rot fungi. The fungus managed to survive in high concentrations of the organo pollutants and degraded 90% anthracene and 97% nitrobenzene [4].

Fungi are also capable of influencing metal transformation in several types of media such as industrial wastes, low-grade ores and metal bearing minerals Bioaccumulation of metals, such as cadmium, zinc, and cesium, by several fungi has been shown. Not only does the fungal mycelium take up metals they can also transport the metals to the fruiting bodies [1,5].

Although groundwater generally is less likely to be contaminated than surface water, it nonetheless may require contaminant removal. Possible groundwater contaminants include naturally occurring inorganic chemicals that have found their way into the aquifer. Groundwater near the land surface usually teems with microbial life. Bacteria are far more numerous than any other organism in the soil and groundwater.

Molds and other fungi also are common in groundwater near the land surface, where plenty of oxygen is present. Most fungi feed on dead or decaying material. Similar to the bacteria, the numbers of fungi decrease with depth.

The aim of the present study is to isolate fungal diversity associated with well bodies and groundwater.

Material and Methods

Sediment samples were collected from different depths of the well samples has been gathered from the well of marsh area after drying by oven under 50 C°.

The following culture media were used for growth of isolated fungi from sediments:

1. Potato carrot Agar- mixture of potato (20gm), Carrot (20gm) and Agar (20gm) in 1litter of distil water, sterilize by autoclave.
2. Malt Extract Agar-mixture of malt extract (15gm) and agar (20gm) in 1 litter of distil water, sterilize by autoclave.

Article Information**DOI:** 10.31021/jwt.20181114**Article Type:** Research Article**Journal Type:** Open Access**Volume:** 1 **Issue:** 3**Manuscript ID:** JWT-1-114**Publisher:** Boffin Access Limited**Received Date:** May 08, 2018**Accepted Date:** May 28, 2018**Published Date:** May 30, 2018***Corresponding author:****Hussein Al-Nasrawi**

Assistant professor

Southern Technical University

Iraq

Tel no: 9647727881028

E-mail: halnasrawia@yahoo.com

Citation: Hussein Al-Nasrawi. Mycoflora of Ground Water associated with Al-Teeb well in Iraq. J Water Technol Treat Methods. 2018 May;1(3):114**Copyright:** © 2018 Hussein Al-Nasrawi. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 international License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

The present study used dilution technique method to isolate fungi from sediments, which summarize by adding 1 gm of dry sediment to 99 ml of sterilized distil water in a conical flask with shaking for 10 min and drain 1 ml of the suspension by a sterilized pipette, follow by pouring culture medium on the sample in the Petri dish.

Isolates of fungi were identified by the author of present research according to fungal keys published in the following papers [6-17].

Results

43 fungal species were isolated from sediments of Al-Teeb well in marsh area of Iraq during 2017; the isolated species belongs to *Hyphomycetes* 33 species, *Ascomycetes* 6 species and *Zygomycetes* 4 species. As shown in Table 1.

New Record Fungus

One new record fungus belong to *Hyphomycetes* was isolated from sediments of we the well.

Table 1: Fungi isolated from sediments of Al-Teeb well.

No.	Fungi
1	<i>Absidia corymbifera</i> (Cohn) Sacc.& Trotter
2	<i>Aphanoascus fulvescens</i> (Cooke)Apinis
3	<i>Alternaria alternata</i> Keissler
4	<i>Alternaria raphani</i> Groves & Skolko
5	<i>Alternaria</i> sp.
6	<i>Allysedium</i> sp.
7	<i>Aspergillus flavus</i> Link & Fries
8	<i>A.fumigatus</i> Fres
9	<i>A.nidulans</i> (Eidam) vuill
10	<i>A.niger</i> van Tieghem
11	<i>A.terreus</i> Thom
12	<i>Aspergillus</i> sp.
13	<i>Aureobasidium pullulans</i> (De Bary) Arnaud
14	<i>Bipolaris hawaiiensis</i> (M.B.Ellis) Subram.&Jain
15	<i>Bipolaris spicifera</i> (Bain)v.Arxx
16	<i>Chaetomium globosum</i> Kunze & Fries
17	<i>Chaetomium</i> sp.
18	<i>Chuppia sarcinifera</i> Deighton
19	<i>Cladosporium cladosporoides</i> (Fresen) de Vries
20	<i>Cladosporium spongiosum</i> Berk.& Curt.
21	<i>Cladosporium</i> sp.
23	<i>Curvularia penniseti</i> (Metra)Boedijn
24	<i>Emericella nidulans</i> (Eidam) vuill .
25	<i>Emericella</i> sp.
26	<i>Eurotium</i> sp.
27	<i>Fusarium oxysporum</i> Schlecht.
28	<i>Fusarium</i> sp.
29	<i>Graphium putredinis</i> (Corda)Hughes
30	<i>Monodictys glauca</i> (Cooke & Harken)Hughes
31	<i>Mucor</i> sp.
32	<i>Penicillium</i> sp.
33	<i>Phoma glomerata</i> (Corda)Wollen.& Hochapfel
34	<i>Phoma herbarum</i> Westend.
35	<i>Phoma</i> sp.
36	<i>Rhizopus stolonifer</i> (Ehrenb:Fr.)Vuill.
37	<i>Stachybotrys atra</i> Corda
38	<i>Stachybotrys atra</i> Corda var. <i>microspora</i> Mathur& Sankhla
39	<i>Stachybotrys</i> sp.
40	<i>Stemphylium</i> sp.
41	<i>Torula herbarum</i> (Pers.)Link ex S.F.Gray
42	<i>Trichocladium opacum</i> (Corda)Hughes
43	<i>Ulocladium botrytis</i> Preuss

Description of *Allysidium* sp. as a new record fungus

Colony black, mycelium partly superficial, partly immersed, hypha branched, septate, dark brown to blackish smooth, 6-8 μ thick, conidiophores with 1-2 branched, semimacronematous, pale to dark colored, conidiogenous cell monoblastic, conidia in branched chains, spherical to oval, or limoniform, 6-9 μ in diameter, without septa fungus isolated from sediment was kept in Basrah university herbarium under the number BSRA 3001 (Figure 1).




Figure 1: Picture of *Allysidium* sp.

Discussion

Aquatic fungi groups *Hyphomycetes*, *Ascomycetes* and *Zygomycetes* primarily originate in water by transferring of pathogenic fungi associated with terrestrial plants to aquatic habitats and bears reproductive units of fungi (spores) during a period of time to adapt with the new habitat. Sediments enriched with nutrients especially organic matter which encourage microorganisms growth [18].

Fungi can survive under sever conditions and has the ability to grow under limited conditions to produce a hypha and to sporulate again or produce fruit bodies as in *Ascomycetes* and *Basidiomycetes*. As shown in Table 1 many species of fungi were isolated from sediment of Al-Teeb well due to suitable conditions of fungal growth specially biodegradation role of fungi in decomposition process, which led to high water purification quality without using chlorination technique. Biological parameters of raw water of the well approached to international standard specification of *E. coli* per 100 ml water. As shown in table 1 many species of fungi were isolated from well sediment.

References

- Gadd GM, Ramsay L, Crawford JW, Ritz K. Nutritional influence on fungal colony growth and biomass distribution in response to toxic metals. *FEMS Microbiol Lett.* 2001 Nov; 204(2):311-316.
- Anand B, Harit J. Degradation of alkali lignin by two ascomycetes and free radical scavenging activity of the products. *Journal Biocatalysis and Biotransformation.* 2017 May;35(4):226-231.
- Helena MB, Oliveira CS, Russell M, Paterson NB, Nelson L. Fungi from a Groundwater-Fed Drinking Water Supply System in Brazil. *Int J Environ Res Public Health.* 2016 Mar;13(3):304.
- Rodriguez C. Industrial and environmental applications of white-rot fungi. *Mycosphere.* 2017 Jan;8(3):456-466.
- Gadd GM. Interaction of fungi with toxic metals *New phytol.* 1993;124:25-60.
- Guarro JA, Figueras MJ. The ascomycetes genus *Chaetomium*, *Novahydwigia* J.Cramer, Berlin. 1986.
- Arx JA, Figueras MJ, Guarro J. Sordaraceous Ascomycetes without Ascospores ejaculation. *Beithelte Zur, Nova Hedwigia.* 1988; 94:1-104.

8. Boerema GH. Contributions towards a monograph of Phoma (Coelomycetes)-II, section Pyrenomyces. Personia. 1993; 15: 197-221.
9. Booth C. The genus Fusarium. Common wealth Mycological Institute. Kew, Surrey, England. 1971;237.
10. Domsch KH, Gams W, Enderson T. Compenedium of soil fungi. Academic press. 1980;1:859.
11. Ellis MB. Dematiaceous Hyphomycetes. Commonwealth Mycological Institute. Kew, Surrey, England. 1971;608.
12. Ellis MB. More dematiaceous Hyphomycetes. Common wealth Mycological Institute .Kew, Surrey, England. 1976;507.
13. Goh TK, Clement KMT. Key to common dematiaceous hyphomycetes from freshwater. Fungal diversity Research series. 2003; 10:325-343.
14. Hoog GS, Guarro J. Atlas of clinical fungi. Gentra albureau Voor Schimmel cultures. Universitat Rovera in Virgili, Spain. 1995; 720.
15. Kohlmeyer J, Kohlmeyer E. Marine Mycology. The Higher Fungi. Institute of Marine Sciences, University of North Carolina at Chapel Hill. Morehead City. North Carolina. 1979; 690.
16. Kohlmeyer J, Kohlmeyer V. Illustrated key to the filamentous higher marine fungi. Botanica Marina, 1961; 34:1-61.
17. Pitt JI. The genus penicillium and its teleomorphic states. Academic press, London, Newyork, Torento, Sydney, San Francisco. 1981;634.
18. Garraway MO, Evans RC. Fungal nutrition and physiology. A Wiley-Interscience Publication. Jhon Wiley and Sons. 1984; 401.